

Accuracy and Equity in Autism Screening, Identification and Referral

Date:

May 2020

Download Material

Autism spectrum disorder (ASD) now affects at least 1 in 59 children. While we know that we can often detect ASD by age 2, the average age of diagnosis remains 4 years old and is even later for racial/ethnic minority, rural and low-income youth and girls. This is concerning as early diagnosis can lead to earlier intervention, which is known to significantly improve outcomes for a child's health and well-being.

In this Research at a Glance brief, we review two PolicyLab and Children's Hospital of Philadelphia studies that provide insight into the reality of screening for ASD and the need to address disparities in screening and referrals to ensure that children with ASD can get the care they need as early as possible.

Authors:

Wallis KE, Guthrie W

Topics

Equitable Access to Services
Early Childhood

Related Content

Closing the Gap: Creating More Equity in Diagnosis of Autism Spectrum Disorder

Adherence to Screening and Referral Guidelines for Autism Spectrum Disorder in Toddlers in Pediatric Primary Care

<u>Screening Instruments for Developmental and Behavioral Concerns in Pediatric Hispanic Populations in the United States: A Systematic Literature Review</u>

Accuracy of Autism Screening in a Large Pediatric Network

<u>Distinguishing Primary and Secondary Early Intervention Programs: Implications for Families, Clinicians and Policymakers</u>

Identifying Autism Spectrum Disorder in Real-world Health Care Settings

Provider Responses to Positive Developmental Screening: Disparities in Referral Practices?

Insights from Behavioral Economics: A Case of Delayed Diagnosis of Autism Spectrum Disorder

USPSTF Request for Public Comment: Screening for Autism Spectrum Disorder in Young Children

No More Watchful Waiting for Developmental Delays

<u>Disparities in Autism Screening and Well-Child Care Persisted During the Pandemic, but Novel Solutions to Promote Equity Exist</u>