

## Simulation of Contraceptive Access for Adolescents and Young Adults Using a Pharmacist-Staffed e-Platform: Development, Usability, and Pilot Testing Study

Date:

Feb 2025 Visit Article

**Background**: Offering contraceptive methods at pharmacies without a prescription is an innovative solution to reduce the incidence of unintended pregnancies among adolescents and young adults (AYA). Pharmacy-prescribed contraception may increase the convenience, simplicity, and affordability of contraceptives.

**Objective**: The aim of this study was to develop, pilot test, and evaluate the acceptability and feasibility of a telemedicine electronic platform app simulating pharmacist prescribing of contraceptives to AYA as well as assess agreement between pharmacist-simulated contraceptive approvals and contraception as prescribed in routine clinic visits.

**Methods**: This study was conducted in two phases: (1) development and usability testing of a prototype app to simulate pharmacists prescribing contraceptives to AYA and (2) pilot testing the app in a simulation for AYA requesting contraception from a pharmacist with pharmacist review and request approval or rejection. Eligibility criteria in both phases included the following: assigned female sex at birth, age 15-21 years, seeking contraceptive services at an academic adolescent medicine clinic, prior history of or intention to have penile-vaginal intercourse in the next 12 months, smartphone ownership, and English language proficiency. Phase 1 (usability) involved a video-recorded "think aloud" interview to share feedback and technical issues while using the app prototype on a smartphone and the completion of sociodemographic, sexual history, and perception of the prototype surveys to further develop the app. Phase 2 (pilot) participants completed phase 1 surveys, tested the updated app in a simulation, and shared their experiences in an audio-recorded interview. Descriptive analyses were conducted for quantitative survey data, and thematic analyses were used for interview transcripts.

**Results**: Of the 22 participants, 10 completed usability testing, with a mean age of 16.9 (SD 1.97) years, and 12 completed pilot testing, with a mean age of 18.25 (SD 1.48) years. Three issues with the prototype were identified during "think aloud" interviews: challenges in comprehension of medical language, prototype glitches, and graphic design suggestions for engagement. Usability testing guided the frontend and backend creation of the platform. Overall, participants agreed or strongly agreed that using an app to receive contraceptives would make it easier for teens to access (n=19, 86%) and make contraceptive use less stigmatizing (n=19, 86%). In addition, participants agreed that receiving contraception prescriptions from a pharmacist without a clinic visit would be safe (n=18, 82%), convenient (n=19, 86%), acceptable (n=18, 82%), and easy (n=18, 82%). Pharmacists and medical providers had 100% agreement on the prescribed contraceptive method for pilot participants.

**Conclusions**: AYA found contraceptive prescription by a pharmacist via an app to be highly acceptable and provided critical feedback to improve the design and delivery of the app. Additionally, pharmacist contraceptive approvals and contraception as prescribed in routine clinic visits were identical.

## Journal:

JMIR Pediatrics and Parenting Authors: Knowles K, Lee S, Yapalater S, Taylor M, Akers AY, Wood S, Dowshen N

## Topics

Sexual & Reproductive Health