

# How Can HPV Vaccination Rates be Improved?

## Statement of Problem


Human papilloma virus (HPV) is most commonly known to cause genital warts, but may also lead to cervical cancer, a significant cause of mortality in women. Despite clinical trials showing the HPV vaccine is safe and effective, vaccination rates remain low among adolescent girls (35 percent fully vaccinated, compared to 70-80 percent for other common adolescent vaccines). Parents report delaying HPV vaccination due to concerns about safety and fears that vaccination may lead to an earlier onset of sexual activity. Additionally, clinician beliefs and practice styles and adolescents' infrequent health care visits may limit receipt of the HPV vaccine, especially since all three doses are required for full protection. We need innovative interventions to overcome these obstacles to vaccine receipt.

Through our research, we aim to inform programs and policy changes that ensure these adolescents can receive care to meet their unique needs and, ultimately, transition to healthy, productive adults. For example, we're investigating how using electronic health records may promote efficient workflows and provide automated support to families and clinicians during health decision-making.

## Description


### How can HPV vaccination rates be improved?

Image



Image

## Percentage Of Vaccinated Girls



Only **35%** of adolescent girls are fully vaccinated against HPV, compared to **70-80%** for other common adolescent vaccines.

PolicyLab developed a three-part clinician-focused decision support intervention (education, electronic health record-based alerts, and feedback), and a family-focused decision support intervention (automated reminder phone calls that also directed families to an educational website). In a one-year clinical trial, 22 primary care practices within Children's Hospital of Philadelphia's Pediatric Research Consortium were randomized to receive clinician-focused decision support or not. Within these practices, 22,486 girls ages 11-17 who were due for the HPV vaccine dose 1, 2, or 3 were randomized to receive family-focused decision support or not. This resulted in four study groups: clinician and family-focused decision support, clinician-focused only, family-focused only and no support. We measured effectiveness of the decision support by comparing final vaccination rates and time to vaccination between the study groups.

Decision support directed at both clinicians and families most effectively increased HPV vaccination rates. Clinician-focused reminders, education, and feedback were most effective for promoting the initiation of the HPV vaccine series, while the family-focused reminder phone calls supported completion of the vaccine series. Both types of decision support (family-focused and clinician-focused) accelerated vaccination uptake. The cost of implementing the most effective single intervention for each dose was low ( $\leq \$10$  per additional girl vaccinated).

### Next Steps

To increase HPV vaccination rates, it is most effective to engage both the family and the clinician. These results demonstrate how health information technology can be utilized to engage both clinicians and families in order to improve health and health care outcomes. PolicyLab has disseminated these findings both locally and nationally, seeking to assist other healthcare providers in boosting their use of the HPV vaccine.

*This project page was last updated in July 2019.*

### Suggested Citation

Children's Hospital of Philadelphia, PolicyLab. *How Can HPV Vaccine Rates be Improved?* [Online] Available at: <http://www.policylab.chop.edu> [Accessed: plug in date accessed here].

## PolicyLab Leads



[Alexander Fiks](#)

MD, MSCE

Faculty Member

## Team



[Kristen Feemster](#)

MD, MPH, MSHP

Faculty Scholar



[Stephanie Mayne](#)

PhD, MHS

Faculty Member

Lihai Song, MS

Robert W. Grundmeier

Dean Karavite

Cayce C. Hughes

James Massey

Ron Keren

Louis M. Bell

Richard Wasserman

Russell Localio

## Funders of Project

## Project Contact

Stephanie Mayne

[MayneS@email.chop.edu](mailto:MayneS@email.chop.edu)

## Related Tools & Publications

- [Improving HPV Vaccination Rates Tools and Memos](#)  
Mar 2014
- [Sociodemographic Differences in Human Papillomavirus Vaccine Initiation by Adolescent Males Article](#)  
Sep 2015
- [Improving HPV Vaccination Rates Using Maintenance-of-Certification Requirements Article](#)  
Feb 2016
- [Increasing HPV Vaccination Rates Among Adolescents: Challenges and Opportunities Evidence to Action Briefs](#)  
Apr 2016
- [HPV Vaccine Recommendation Profiles Among a National Network of Pediatric Practitioners: Understanding Contributors to Parental Vaccine Hesitancy and Acceptance Article](#)  
Dec 2018
- [Disparities in HPV Vaccine Series Completion by Adolescent Males: A Retrospective Cohort Study Article](#)  
May 2019
- [Missed Opportunities for Human Papillomavirus Vaccination at Office Visits During Which Influenza Vaccine Was Administered: An AAP Pediatric Research in Office Settings \(PROS\) National Primary Care Research Network Study Article](#)  
Jul 2020
- [HPV Vaccinations at Acute Visits and Subsequent Adolescent Preventive Visits Article](#)  
Nov 2022
- [Provider Perspectives on Clinical Decision Support to Improve HIV Prevention in Pediatric Primary Care: A Multiple Methods Study Article](#)  
Feb 2023
- [What Health Systems Can Do Now to Improve Human Papillomavirus Vaccination Article](#)  
Nov 2023
- [Missed Opportunities for Adolescent Immunizations at Well-Care Visits During the COVID-19 Pandemic Article](#)

